# Upper Ocean Dynamics in the Tropical and South Atlantic from High Density XBT Lines AXXII and AX18

Gustavo J. Goni, Molly O. Baringer, Silvia Garzoli,

Qi Yao and Claudia Schmid

**United States Department of Commerce** 

**National Oceanic and Atmospheric Administration** 


Atlantic Oceanographic and Meteorological Laboratory

Miami, Florida, USA.

Gustavo.Goni@noaa.gov


#### **Motivation**


\* 2(AX18) 4(AX08) Times a year \* Every 40 km




- Investigation of the variability of zonal currents in the tropical Atlantic to aid in the study of the role of upper ocean dynamics on sea surface temperature.
- Meridional water mass transfer across the equator.
- Investigation of the meridional heat transport across 30-35S.

# Historical XBT data along AX08









Figures by R. Molinari


# **AX08** and the major surface currents in the tropical Atlantic




# Sea height variability




## The first six transects in AX08




# Sea height anomaly field




### Temperature section





### Dynamic height





### Cumulative geostrophic transport














-20C







## Profiling float deployment on AX08

- 15 profiling floats deployed in Dec 2000
- 40+ profiling floats deployed to this date





#### Salinity uncertainties & Dyn Heights





Will soon be improved, expanded

Current	December 2000		September 2001		January 2002	
	location	Sv	location	Sv	location	Sv
NEC	19.7-9.7°N	-23	19-13.1°N	-15	19.8-11.1°N	-22
NECC	7.2-4.9°N	25	9.7-5.6°N	22	9.3-7°N	16
nSEC	4.9-3.7°N	-20	5.6-3.2°N	-20	7-5.5°N	-14
NEUC	3.7-3.0°N	14	-	-	5.5-3.9°N	18
nSEC	3.0-1.7°N	-37	-	-	3.9-2.2°N	-30
eSEC	1.5-3.6°S	-36	1.7-4.3°S	-5	1.8-4.2°S	-13
SEUC	3.6-5.2°S	17	4.3-5.4°S	11	4.2-4.7°S	14
cSEC	5.2-6.8°S	-13	5.4-6.9°S	-11	4.7-8.0°S	-6
SECC	10-12.9°S	6	6.7-9.5°S	11	8.0-9.3°S	7
sSEC	12.9-19.9°S	-7	9.5-15.3°S	-10	9.3-20°S	-8


#### **IMPROVEMENTS**

- Intermediate to high density XBT deployment to Cape Town, underway
- •Include TSG, too problematic, expensive
- •Include XCTD, expensive, not included in current budget
- •Include deeper XBTs, to 2000 m deep, expensive, redesign of autolaunchers, not included in current budget
- •More profiling float data, depending on funding and interest


#### Current and Immediate future work

- \* Obtain mean locations, transports, from monthly mean temperature sections.
- \* Relationship between surface currents and sea height signature.
- \* Investigate and (maybe) quantify the effect of waves in the synoptic temperature sections, the dynamic height and transport estimates.
- \* Compare our results with estimates from drifters and CM in PIRATA mooring (33°W).
- \* Compare our results with estimates obtained from numerical models.
- \* Use these results to improve the methodology used to monitor surface currents in NRT from altimetry in CoastWatch.


### The first three transects of AX18





# The transects and the upper ocean dynamics



#### **Cumulative geostrophic transports**







ring C

20

Agulhas ring A


20°E

10°E

# **AX18** and the Meridional heat flux in the Atlantic Ocean

In the South Atlantic the meridional heat flux is uncertain, estimates vary between -0.4 and +0.9 PW.


To better understand the global ocean thermohaline circulation and its impact on climate it is necessary to reduce the heat flux uncertainty in the South Atlantic



# Estimates of South Atlantic meridional heat flux near 30°S


Lat °S	Heat Flux PW	Method /	Source
<b>32</b>	0.66-0.88	Inverse	Fu (1981)
30	0.69	Sea-air fluxes	Hastenrath (1982)
<b>32</b>	0.16-0.68	Direct	Bennett (1978)
32.5	0.63	Numerical model	Donners
<b>32</b>	0.4	Direct	Bryan (1962)
30	0.39	Sea-air fluxes	Bunker (1980)
30	0.38	Sea-air fluxes	Hsiung (1985)
<b>30</b>	0.3	Inverse	Macdonald & Wunsch (1996)
			Ganachaud & Wunsch (2000)
30	0.29	Numerical model	Marchesiello et al. (1998)
30	0.26	Numerical model	Matano (pers. comm., 2003)
<b>32</b>	0.24	Inverse	Rintoul (1991)
<b>30</b>	0.22	Direct	McDonogh and King (2003)
30	0.19	Numerical model	Matano & Philander (1993)
30	- 0.23	Inverse	de las Heras & Schlitzer (1999)

# The first two estimate of meridional heat flux



#### Data distribution

www.aoml.noaa.gov/phod/hdenxbt




#### **Data distribution**

**AX08** 

Proposal (PDF format)

**AX18** 



#### **Current and Immediate future work**

- Validate altimeter estimates of the transport of the Brazil and Agulhas currents.
- Validate altimeter estimates of the motion of the Brazil Current front.
- Continue with the estimates of meridional heat transport across 35°S.